



Near Earth Object Observations Program

Presentation to
**UN COPUOS
Scientific & Technical
Subcommittee**

Lindley Johnson
Program Executive
NASA HQ
16 February 2010



Terminology



- “Near Earth Objects (NEOs)”- any small body (comet or asteroid) passing within 1.3 Astronomical Unit (AU) of the Sun
 - 1 AU is the distance from Earth to Sun = ~ 150 million kilometers (km)
 - NEOs are predicted to pass within ~ 45 million km of Earth’s orbit
 - Population of:
 - Near Earth Asteroids (NEAs)
 - Near Earth Comets (NECs) – also called Earth Approaching Comets (EACs)
 - 84 currently known
- “Potentially Hazardous Objects (PHOs)” – small body that has potential risk of impacting the Earth at some point in the future
 - NEOs passing within 0.05 AU of Earth’s orbit
 - ~ 8 million km = 20 times the distance to the Moon
 - Appears to be about 20% of all NEOs discovered



NEO Observation Program



US component to International Spaceguard Survey effort
Has provided 98% of new detections of NEOs

Began with NASA commitment to US Congress in May, 1998

Scientific Objective: Discover 90% of NEOs larger than 1 kilometer in size within 10 years (1998 – 2008)

NASA Authorization Act of 2005 provided additional direction (but no additional funding)

“ . . . plan, develop, and implement a Near-Earth Object Survey program to detect, track, catalogue, and characterize the physical characteristics of near-Earth objects equal to or greater than **140 meters** in diameter in order to assess the threat of such near-Earth objects to the Earth. It shall be the goal of the Survey program to achieve **90 percent completion** of its near-Earth object catalogue **within 15 years** [by 2020].



NASA's NEO Search Program

(Current Systems)





NEO Program Office @ JPL

- Program coordination
 - Automated SENTRY
- www.neo.jpl.nasa.gov

Minor Planet Center (MPC)

- IAU sanctioned
 - Discovery Clearinghouse
 - Initial Orbit Determination
- www.cfa.harvard.edu/iau/mpc.html






Minor Planet Center Upgrade



- MPC fully upgraded to LINUX-based processors
- Processes all observations received worldwide each night within next day
- Database contains:
 - Over 68,000,000 observations
 - Over 475,000 orbits for minor planets
- NEO observations identified and processed on receipt in near-real-time
- Suspected discoveries automatically posted to NEO Confirmation Page
- Adequately prepared for next generation search systems

SKY COVERAGE

Plot prepared 17 Nov 2009 14:17:18 by the Minor Planet Center



2009/10/01

2009/11/01

Morning

Evening

12^h

08^h

04^h

00^h

20^h


16^h

Opposition Point = 01 25.6,+08 59. Fields reaching fainter than $V = 18.0$.

- | | | | | |
|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|
| 2009/11/01 (2009 305) | 2009/10/31 (2009 304) | 2009/10/30 (2009 303) | 2009/10/29 (2009 302) | 2009/10/28 (2009 301) |
| 2009/10/27 (2009 300) | 2009/10/26 (2009 299) | 2009/10/25 (2009 298) | 2009/10/24 (2009 297) | 2009/10/23 (2009 296) |
| 2009/10/22 (2009 295) | 2009/10/21 (2009 294) | 2009/10/20 (2009 293) | 2009/10/19 (2009 292) | 2009/10/18 (2009 291) |
| 2009/10/17 (2009 290) | 2009/10/16 (2009 289) | 2009/10/15 (2009 288) | 2009/10/14 (2009 287) | 2009/10/13 (2009 286) |
| 2009/10/12 (2009 285) | 2009/10/11 (2009 284) | 2009/10/10 (2009 283) | 2009/10/09 (2009 282) | 2009/10/08 (2009 281) |
| 2009/10/07 (2009 280) | 2009/10/06 (2009 279) | 2009/10/05 (2009 278) | 2009/10/04 (2009 277) | 2009/10/03 (2009 276) |




WISE Finds First NEO



First NEO discovered by WISE: 2010 AB78

- Diameter ~ 600 m
- Amor (Mars crosser)
- Not a PHO
- Observed by WISE
1/12/10, before sky
survey start






Discovery Metrics

Discovery Rate of >1km NEOs



Large NEO Discovery Rate






Discovery Metrics




Cumulative Large NEO Discoveries






Known Near Earth Asteroid Population






Known Near Earth Asteroid Population






Update on Apophis



Predicted Close Approach
of 2004 MN4 “Apophis”
(a ~270 meter object)
on April 13, 2029



With improved processing of acquired observations the probability of threat on the subsequent 2036 pass has dropped from 1 in 43,000 to less than 1 in 250,000



NRC NEO Report*

Findings & Recommendations (1 of 2)



SURVEY and DETECTION:

Finding: The current Near-Earth Object surveys cannot meet the goal to discover 90 percent of all NEOs 140 meters in diameter or greater by 2020.

Finding: The optimal approach to achieving this NEO Survey goal will depend on nonscientific factors. If time to complete should be minimized, a space-based capability augmented by large aperture ground observatories is most effective. If cost should be minimized, large aperture ground observatories acquired for other purposes could complete survey in under 20 years

Recommendation: Recent studies suggest that objects as small as 30 to 50 meters in size could be highly destructive. Surveys should attempt to detect as many 30- to 50-meter objects as possible.

CHARACTERIZATION:

Finding: The Arecibo and Goldstone planetary radars play a unique role in the characterization of NEOs, providing unmatched accuracy in orbit determination, and insight into size, shape, surface structure

Recommendation: Immediate action is required to ensure the continued operation of the Arecibo Observatory at a level sufficient to maintain and staff the radar facility.

*National Research Council : “Defending Planet Earth: Near Earth ¹³
Object Surveys and Hazard Mitigation Strategies”, Jan 22, 2010





Radar Studies



- Observations on the limited accessible objects
- 20 to 30 NEOs/year from Goldstone and Arecibo
 - Required for timely precision orbit determination
 - Characterization with sufficient signal strength
 - Shape, spin-state, surface structure
 - Satellites (and then derived mass)



Study of Shape, Size, Motion and Mass of 66391 (1999 KW4)



Shape, Size of 6489 Golevka





NRC NEO Report*

Findings & Recommendations (2 of 2)



MITIGATION:

Finding: No single approach to mitigation is adequate to fully prevent the effects of the full range of potential impactors, although civil defense is an appropriate component of mitigation in all cases. With adequate warning, a suite of four types of mitigation [Civil Defense, “Slow Push”, Kinetic Impact, Nuclear Detonation] is adequate to mitigate the threat from nearly all NEOs except the most rare energetic ones

Recommendation: The United States should initiate a peer-reviewed, targeted research program in the area of impact hazard and mitigation of NEOs. The scope should include analysis, simulation, and laboratory experiments.

Recommendation: First priority for a space mission in the mitigation area is a test of a kinetic impactor along with a characterization, monitoring and verification system, such as the Don Quijote mission that was considered by ESA. This mission would produce the most significant advances in understanding and provide an ideal chance for international collaboration in a realistic mitigation scenario.

INTERNATIONAL COOPERATION:

Recommendation: The United States should play a significant role in organizing and empowering a suitable international entity to participate in developing a detailed plan for dealing with the NEO hazard.

*National Research Council : “Defending Planet Earth: Near Earth Object Surveys and Hazard Mitigation Strategies”, Jan 22, 2010



Future Years Outlook



-
- Funding has been appropriated to support research with Arecibo planetary radar through 2011
 - As funding becomes available for the purpose, the United States will continue to upgrade and acquire improved NEO detection/characterization capability and begin more focused research on mitigation methods and strategies.